

Characteristic Dynkin Diagrams and W algebras.

E. Ragoucy*

*NORDITA, Blegdamsvej 17,
DK-2100 Copenhagen Ø, Denmark*

Abstract

We present a classification of characteristic Dynkin diagrams for the A_N , B_N , C_N and D_N algebras. This classification is related to the classification of $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras arising from non-Abelian Toda models, and we argue that it can give new insight on the structure of W algebras.

*On leave from absence of Laboratoire de Physique Théorique ENSLAPP, Chemin de Bellevue BP 110, F - 74941 Annecy-le-Vieux Cedex, France. E-mail: RAGOUCY@FRLAPP51.IN2P3.FR

1 Introduction

These recent years, a lot of efforts have been made in order to study the extended conformal algebras [1], also called W algebras. Two of the main problems concerning these algebras are that, first, they have non linear commutation relations, so that there are technical difficulties to deal with, and, second, that there exists no classification scheme, so that a global point of view for their study is hard to find. In order to classify these algebras, one needs to determine which features are relevant to fully distinguish their structure. Of course, the conformal spin contents will be one of them, but it is known that it is not sufficient to define a W algebra. Some results have been obtained in that direction [2], for a wide class of W algebras, namely the so-called $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras that arise in the framework of non-Abelian Toda theories [3].

In this paper, we propose what could be thought of as the Dynkin diagrams (DDs) for $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras: the Characteristic Dynkin Diagrams (CDDs). The CDDs (introduced by Dynkin [4]) are DDs where each node carry a grade 0, $\frac{1}{2}$, or 1, and are in one to one correspondence with the $Sl(2)$ subalgebra of an algebra $\mathcal{G}$ (themselves being known as classifying the $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras). Until now, only the CDDs of exceptional algebras were classified [4], and we present in the first sections of this paper a classification of CDDs for the other algebras. Then, we will connect the CDDs to the $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras and show how they can be of some help in the study of these algebras. Finally, we will discuss some open problems in the conclusion.

2 Some basic notations and definitions

A *Characteristic Dynkin Diagram (CDD)* is built from a Dynkin Diagram (DD) of a simple Lie algebra $\mathcal{G}$ in such a way that it characterizes an $Sl(2)$ subalgebra of $\mathcal{G}$. Practically, to any node in the DD of $\mathcal{G}$, one attaches the grade of the corresponding simple root generator under the $Sl(2)$ -Cartan generator. It can be shown that these grades can only be 0, $\frac{1}{2}$, or 1. However, to any choice of grades (in $\{0, 1, \frac{1}{2}\}$) will not always correspond an $Sl(2)$ algebra, since there exist some grading operators in $\mathcal{G}$ that are not Cartan generator of an $Sl(2)$ $\mathcal{G}$ -subalgebra. In other words, among the 3^n possibilities of constructing a CDD from a rank n algebra, only a certain number of them will really be CDDs. In sections 3,4,5 we present a complete classification of the CDDs for the algebras A_N , B_N , C_N , and D_N . The CDDs of the exceptional algebras have already being classified by Dynkin [4], and we will not study them here.

Let us also recall that the $Sl(2)$ subalgebras of A_N , B_N and C_N are classified by the regular subalgebras of these algebras. In the case of D_N , one has to add the irregular embeddings of the type $B_M \oplus B_{N-M}$ with $2M \neq N$. Still for D_N algebras, some $Sl(2)$ subalgebras have a multiplicity 2 (3 for D_4), which means that there are two (three) non-conjugate $Sl(2)$ related by an outer automorphism of D_N (D_4). Using CDDs, this multiplicity is given by the quotient of the number of symmetries of its CDD by the number of symmetries of the DD: thus, the multiplicity of any $Sl(2)$ will come naturally with its CDD, and we do not have to pay attention to it.

We will use intensively the connection that exists between CDDs and defining vectors. A defining vector $f = (f_1, f_2, \dots, f_N)$ is formed by the components f_i of the $Sl(2)$ -Cartan generator in a given basis. For a good choice of basis, the grades of the simple roots (i.e. the nodes of the CDD) can be expressed in a simple way as function of the f_i 's: for any algebra $\mathcal{G}=A, B, C$ or D , and denoting by e_i a basis of $\mathfrak{R}^N$ where $N=\text{rank}\mathcal{G}$ ($\text{rank}\mathcal{G}+1$ for A_M), the grade of simple roots

of type $e_{i+1} \pm e_i$ (when they exist in $\mathcal{G}$) will be $f_{i+1} \pm f_i$ and the grade of $2e_i$ and e_i will be $2f_i$ and f_i (respectively). With this normalisation, the defining vector of any $Sl(2)$ can be directly read from its CDD, and vice versa.

Finally, let us introduce some notations:

We will call *integral CDD*, a CDD where all the coefficients are 0 or 1. The corresponding gradation of the algebra will be integral. Among the integral CDDs, the *principal CDD* is the one where all the nodes have grade 1: it correspond to the principal $Sl(2)$ in $\mathcal{G}$.

A *half-integral CDD* will be a CDD where the coefficients are only 0 or $\frac{1}{2}$, whereas a *mixed CDD* will contain 0, $\frac{1}{2}$ and 1. These two types of CDD correspond to half-integral gradations.

We will call a *piece of length p* any part of a CDD formed by p nodes. Thus, a node is also a piece of length 1.

We will also draw some CDDs as

$$(\overset{s_1}{\bigcirc}-) \overset{s_2}{\bigcirc}-\overset{s_3}{\bigcirc}-\overset{s_4}{\bigcirc}-\dots \dots -\overset{s_n}{\bigcirc} \quad (2.1)$$

where the notation $(\overset{s_1}{\bigcirc}-)$ means that both CDDs with and without the node $\overset{s_1}{\bigcirc}$ have to be considered. For this kind of CDDs, we will use a plus (+) and a minus (-) sign to distinguish the CDD with the node from the other (without the node). In the above example, these CDDs will be denoted as (2.1^+) and (2.1^-) .

3 Characteristic Dynkin Diagrams for A_N

Let us first remark that the CDDs of A_N are symmetric. This can be proven in two ways. First, because the $Sl(2)$ -Cartan generator can be chosen diagonal in A_N : as it is traceless, it can be chosen symmetric, so that the corresponding CDD is also symmetric. Second, because any $Sl(2)$ has a multiplicity one: since the DD of A_N possesses a $\mathbb{Z}_2$ symmetry, so must any CDD. Notice that the symmetry centre of A_N is either a node (for A_{2N+1}), or a link between two nodes (for A_{2N}). We will indicate the middle (i.e. the symmetry centre) by a double arrow $\updownarrow$, and draw only half of each CDD of A_N .

3.1 Integral CDDs of A_N

Let us look first at an integral CDD. It is of the form

$$\underbrace{\overset{1}{\bigcirc}-\dots-\overset{1}{\bigcirc}}_{n_1} \underbrace{\overset{0}{\bigcirc}-\dots-\overset{0}{\bigcirc}}_{p_1} \underbrace{\overset{1}{\bigcirc}-\dots-\overset{1}{\bigcirc}}_{n_2} \underbrace{\overset{0}{\bigcirc}-\dots-\overset{0}{\bigcirc}}_{p_2} \dots \dots \underbrace{\overset{1}{\bigcirc}-\dots-\overset{1}{\bigcirc}}_{n_1} \quad (3.1)$$

where we can suppose $p_i \neq 0$ ($\forall i$) without any loss of generality. The gradation H associated to this CDD will be of the form


$$(x, x-1, x-2, \dots, x-n_1, \underbrace{x-n_1, \dots, x-n_1}_{p_1}, x-n_1-1, \dots, x-n_1-n_2, \underbrace{x-n_1-n_2, \dots, x-n_1-n_2}_{p_2}, \dots) \quad (3.2)$$

with x determined by $tr H = 0$. H is an $Sl(2)$ -Cartan generator, so that its eigenvalues can be gathered in $\mathcal{D}_j$ representations. This implies that for any eigenvalue y in (3.2), $y-1$ must also


appear with a multiplicity which is greater than or equal to the multiplicity of y . For $y = x - n_1$, this means $x - n_1 - 1$ must appear at least $(p_1 + 1)$ times, so that $n_2 = 1$ and $p_2 \geq p_1$. The same argument for all the eigenvalues $y_j = x - \sum_{i=1}^j n_i$ leads us to the constraints:

$$n_i = 1, \text{ and } p_i \geq p_{i-1} \quad \forall i \geq 2 \quad (3.3)$$

Then, the symmetry of the CDD will lead to the two types of integral CDDs for A_N :




$$p_1 \leq p_2 \leq \dots \leq p_\ell \quad (3.4)$$



$$p_1 \leq p_2 \leq \dots \leq p_\ell \quad (3.5)$$

3.2 Half-integral CDDs of A_N

For half-integral CDDs, the above reasoning is still valid, with the difference that between the eigenvalues y and $y - 1$, there will occur the eigenvalue $y - \frac{1}{2}$, which is in a different $Sl(2)$ representation. Instead of one series of p_i 's (as in the integral case), we obtain two independent series (p_i and q_i) which are intertwined. Thus, the half-integral CDDs take the form:




$$p_1 \leq p_2 \leq \dots \leq p_\ell \text{ and } q_1 \leq q_2 \leq \dots \leq q_{\ell'} ; u = p_\ell \text{ or } q_{\ell'} \text{ depending on } \ell' = \ell - 1 \text{ or } \ell \quad (3.6)$$

Notice that the CDD (3.6) can be split in two different CDDs, depending on the two different values $(\ell - 1)$ and ℓ that ℓ' can take.

3.3 Mixed CDDs for A_N

The computation seems very much more complicated when looking at the mixed CDDs. However, it can be greatly simplified if one remarks that the pieces of CDDs that contains nodes of grade 1 must be outside of the pieces containing nodes with grade $\frac{1}{2}$. In fact, looking at a piece of CDD which possesses a grade 1 node inside two grade $\frac{1}{2}$ nodes, one computes:



$$\longrightarrow H = (\dots, x, x - \frac{1}{2}, \dots, x - \frac{1}{2}, x - \frac{3}{2}, \dots, x - \frac{3}{2}, x - 2, \dots)$$

$\underbrace{\hspace{10em}}_{p+1} \quad \underbrace{\hspace{10em}}_{q+1}$

so that the eigenvalue $x - 1$ does not appear, as it should do. Thus, the mixed CDDs are formed "outside" by an integral CDD, and "inside" by a half-integral CDD. To study the "transition" between these two pieces of the mixed CDD is as easy as for the preceding cases. Altogether, the mixed CDDs of A_N take the form:

[illegible]

[illegible]

3.4 $Sl(2)$ decomposition associated to a CDD

In the previous section, we have used the Cartan generator of the $Sl(2)$ subalgebra to determine all the possible CDDs. This generator also allows us to know the $Sl(2)$ representations that enter into the fundamental representation of A_N . A $\mathcal{D}_j$ representation will be characterized by the eigenvalues $(j, j-1, \dots, -j)$ of the Cartan generator. We give in the following table the $Sl(2)$ decomposition for each CDD described in the previous section. To simplify the formulae, we define $p_0 = 0$ and $q_0 = 0$.

CDD	Decomposition of the fundamental of A_{N-1}
(3.4)	$\underline{\mathbb{N}} = \mathcal{D}_{n+\ell-1} \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{\ell-1-i}$
(3.5)	$\underline{\mathbb{N}} = \mathcal{D}_{n+\ell-1/2} \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{\ell-i-1/2}$
(3.6)	$\underline{\mathbb{N}} = \mathcal{D}_{(\ell+\ell'+n-1)/2} \oplus \mathcal{D}_{(\ell+\ell'+n-2)/2} \oplus \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{(\ell+\ell'-1)/2-i} \oplus_{i=0}^{\ell'-1} (q_{i+1} - q_i) \mathcal{D}_{(\ell+\ell'-2)/2-i}$ (3.9)
(3.7) ($n \neq 0$)	$\underline{\mathbb{N}} = \mathcal{D}_{n+(m+\ell+\ell'-1)/2} \oplus \mathcal{D}_{(m+\ell+\ell'-2)/2} \oplus \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{(\ell+\ell'-1)/2-i} \oplus_{i=0}^{\ell'-1} (q_{i+1} - q_i) \mathcal{D}_{(\ell+\ell'-2)/2-i}$
(3.8) ($n + \ell \neq 0$)	$\underline{\mathbb{N}} = \mathcal{D}_{n+\ell+(\ell'+\ell''-1)/2} \oplus_{i=0}^{\ell+\ell'-1} (p_{i+1} - p_i) \mathcal{D}_{\ell+(\ell'+\ell''-1)/2-i} \oplus \oplus_{i=0}^{\ell''-1} (q_{i+1} - q_i) \mathcal{D}_{(\ell'+\ell''-2)/2-i}$

Let us remark that these five decompositions represent all the types of decomposition one can obtain from the reduction of A_N with respect to any of its $Sl(2)$ subalgebras.

4 CDDs of B_N and C_N

The algebras B_N and C_N are irregular subalgebras of A_M ones. A nice way to see how they are

embedded in A_M , is to use the $\mathbb{Z}_2$ symmetry of A_M -DD. This symmetry consists in exchanging the simple root α_i with the simple root α_{M+1-i} . The invariant subalgebra is built on the generators $E_{\alpha_i} + E_{\alpha_{M+1-i}}$, so that the DD of this subalgebra is obtained by folding the A_M -DD. By such a folding, we directly get the B_N and C_N DDs. Here, we will use these foldings $A_{2N} \rightarrow B_N$ and $A_{2N-1} \rightarrow C_N$ to construct not only the DDs, but the CDDs of B_N and C_N from those of A_M . The folding can be extended from the DD to the CDDs of A_M , because the CDDs are themselves symmetric, so that the roots that are summed have the same grade. Of course, only some of the CDDs of A_{2N} (A_{2N-1}) will lead to CDDs of B_N (C_N). The criteria to select them will be the $Sl(2)$ decomposition of the fundamental of A_M , and we will use intensively the table 3.9. As for the A_M case, the integers p_i , q_i and also k_i will satisfy the conditions:

$$\begin{aligned} p_1 \leq p_2 \leq p_3 \leq \dots \leq p_\ell \leq p_{\ell+1} \leq \dots \leq p_{\ell+\ell'} \quad p_0 = 0 \\ q_1 \leq q_2 \leq q_3 \leq \dots \leq q_{\ell'} \text{ or } q_{\ell''} \quad q_0 = 0 \\ k_1 \leq k_2 \leq k_3 \leq \dots \leq k_{\ell+\ell'+\ell''} \quad k_0 = -\frac{1}{2} \quad (\text{so that } 2k_0 + 1 = 0) \end{aligned} \quad (4.1)$$

4.1 CDD for B_N

A CDD of A_{2N} will lead to a CDD of B_N iff all the $\mathcal{D}_j$ with $j \in \frac{1}{2} + \mathbb{Z}$ have an even multiplicity (in the fundamental of A_{2N}). For each decomposition given in table 3.9, we determine which are the cases where the folding can be effectively done, and then give the folded diagram and the corresponding decomposition of the fundamental. Of course, as we are folding A_{2N} , we will consider only the CDDs with an even number of nodes, so that the center of the CDD is always on a link. Notice that under the folding, a CDD of the type (3.6) will behave in two different ways, depending on whether u is equal to or different from 0. If $u = 0$, then the short root of B_N will have a grade $\frac{1}{2}$, while for $u \neq 0$ it will possess a 0-grade.

4.1.1 Integral CDDs

The CDD (3.4) can always be folded in A_{2N} , however, for $p_\ell = 2u + 2 \neq 0$ we get the CDD (4.2) of B_N , while for $p_\ell = 0$ (which implies $p_i = 0 \forall i$), we obtain the principal CDD (4.3). The CDD (3.5) corresponds to A_N with N odd, so that we do not have to consider it here.

$$\begin{array}{c} \begin{array}{cccccccccccccccccccc} \textcircled{1} & \cdots & \textcircled{1} & \textcircled{0} & \cdots & \textcircled{0} & \textcircled{1} & \textcircled{0} & \cdots & \textcircled{0} & \textcircled{1} & \cdots & \textcircled{0} & \cdots & \textcircled{0} & \textcircled{1} & \textcircled{0} & \cdots & \textcircled{0} & \textcircled{0} & \textcircled{0} \\ \underbrace{\hspace{1.5cm}}_n & & \underbrace{\hspace{1.5cm}}_{p_1} & & \underbrace{\hspace{1.5cm}}_{p_2} & & \underbrace{\hspace{1.5cm}}_{p_{\ell-1}} & & \underbrace{\hspace{1.5cm}}_u & & & & & & & & & & & & & \bullet \end{array} \\ \underline{2N+1} = \mathcal{D}_{n+\ell-1} \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{\ell-i-1} \quad p_\ell = 2u + 2 \neq 0 \end{array} \quad (4.2)$$

$$\begin{array}{c} \begin{array}{ccccccc} \textcircled{1} & \textcircled{1} & \cdots & \textcircled{1} & \textcircled{1} & \textcircled{1} & \bullet \end{array} \\ \underline{2N+1} = \mathcal{D}_n \end{array} \quad (4.3)$$

4.1.2 Half-integral CDDs

The CDD (3.6) provides three types of B_N -CDDs, depending on $(n + \ell + \ell' - 1)$ even or odd, $n = 0$ or 1 (only allowed values here) and also¹ $q_\ell = 0$ or $q_\ell > 0$.

(4.4) is obtained from (3.6) with the constraints $n = 0$, $\ell' = \ell$, $p_i = 2k_i + 1$, and $q_\ell = 2u + 2 \neq 0$ for (4.4⁻); and $n = 1$, $\ell' = \ell$, $p_i = 2k_i + 1$, and $q_\ell = 2u + 2 \neq 0$ for (4.4⁺).

¹Note that $q_\ell = 0$ implies $q_i = 0 \forall i$.

The conditions ($n = 0$, $\ell' = \ell$, $p_i = 2k_i + 1$, and $q_\ell = 0$) and ($n = 1$, $\ell' = \ell - 1$, $q_i = 2k_i + 1$, and $p_\ell = 0$) on (3.6) give the CDDs (4.5⁻) and (4.5⁺) respectively.

(4.6) is obtained from (3.6) with the conditions $n = 0$, $\ell' = \ell - 1$, $q_i = 2k_i + 1$, and $p_\ell = 2u + 2 \neq 0$. We recover the CDD (4.5⁺) with the conditions $n = 0$, $\ell' = \ell - 1$, $q_i = 2k_i + 1$ and $p_\ell = 0$ on (3.6).

$$\begin{aligned}
 & \left(\overset{\frac{1}{2}}{\circ} - \right) \underbrace{\overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_1+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{q_1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_2+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{q_2} \cdots \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_\ell+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_u \overset{0}{\bullet} \\
 \frac{2N+1}{q_\ell = 2u+2} = & \begin{cases} \mathcal{D}_{\ell-1/2} \oplus \mathcal{D}_{\ell-1} \oplus_{i=0}^{\ell-1} \left[2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2} \oplus (q_{i+1} - q_i) \mathcal{D}_{\ell-i-1} \right] & \text{without } \overset{\frac{1}{2}}{\circ} \\ \mathcal{D}_\ell \oplus \mathcal{D}_{\ell-1/2} \oplus_{i=0}^{\ell-1} \left[2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2} \oplus (q_{i+1} - q_i) \mathcal{D}_{\ell-i-1} \right] & \text{with } \overset{\frac{1}{2}}{\circ} \end{cases} \quad (4.4)
 \end{aligned}$$

$$\begin{aligned}
 & \left(\overset{\frac{1}{2}}{\circ} - \right) \underbrace{\overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_1+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{\frac{1}{2}}{\circ} \cdots \overset{0}{\circ}}_{2k_2+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{\frac{1}{2}}{\circ} \cdots \overset{0}{\circ}}_{2k_\ell+1} \overset{\frac{1}{2}}{\bullet} \\
 \frac{2N+1}{q_\ell = 2u+2} = & \begin{cases} \mathcal{D}_{\ell-1/2} \oplus \mathcal{D}_{\ell-1} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2} & \text{without } \overset{\frac{1}{2}}{\circ} \\ \mathcal{D}_\ell \oplus \mathcal{D}_{\ell-1/2} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2} & \text{with } \overset{\frac{1}{2}}{\circ} \end{cases} \quad (4.5)
 \end{aligned}$$

$$\begin{aligned}
 & \underbrace{\overset{0}{\circ} \cdots \overset{0}{\circ}}_{p_1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_1+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{p_2} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_2+1} \cdots \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_{\ell-1}+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_u \overset{0}{\bullet} \\
 \frac{2N+1}{p_\ell = 2u+2} = & \mathcal{D}_{\ell-1} \oplus \mathcal{D}_{\ell-3/2} \oplus_{i=0}^{\ell-1} (p_{i+1} - p_i) \mathcal{D}_{\ell-i-1} \oplus_{i=0}^{\ell-2} (k_{i+1} - k_i) \mathcal{D}_{\ell-i-3/2} \quad (4.6)
 \end{aligned}$$

4.1.3 Mixed CDDs

Studying (3.7) and (3.8) leads to the following CDDs:

(4.7) is obtained from (3.7) by the restrictions $m = 1$, $p_i = 2k_i + 1$, $\ell' = \ell$ and $q_\ell = 2u + 2 \neq 0$.

(4.8) with $\ell = 0$ comes from (3.7) by the conditions $m = 0$, $\ell' = \ell - 1$, $q_i = 2k_i + 1$ and $p_\ell = 2u + 2 \neq 0$, while the general case (for (4.8)) is given by (3.8) with $\ell'' = \ell' - 1$, $q_i = 2k_i + 1$ and $p_{\ell+\ell'} = 2u + 2 \neq 0$.

(4.9) is the folding of (3.8) when applying $\ell'' = \ell'$, $n = 0$, $p_i = 2k_i + 1$ and $q_{\ell''} = 2u + 2 \neq 0$.

(4.10) is obtained from (3.8) by the restrictions $\ell'' = \ell'$, $n = 0$, $p_i = 2k_i + 1$ and $q_\ell = 0$.

(4.11) can be obtained in several ways: from (3.7) with the conditions ($\ell' = \ell - 1$, $m = 0$, $q_i = 2k_i + 1$ and $p_\ell = 0$) or ($\ell' = \ell$, $m = 1$, $p_i = 2k_i + 1$ and $q_\ell = 0$), and also from (3.8) when imposing $\ell'' = \ell' - 1$, $q_i = 2k_i + 1$ and $p_{\ell+\ell'} = 0$.

$$\begin{aligned}
 & \underbrace{\overset{1}{\circ} \cdots \overset{1}{\circ}}_n \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_1+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{q_1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_2+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{q_2} \cdots \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_{2k_\ell+1} \underbrace{\overset{\frac{1}{2}}{\circ} \overset{0}{\circ} \cdots \overset{0}{\circ}}_u \overset{0}{\bullet} \\
 \frac{2N+1}{q_\ell = 2u+2} = & \mathcal{D}_{n+\ell} \oplus \mathcal{D}_{\ell-1/2} \oplus_{i=0}^{\ell-1} \left[2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2} \oplus (q_{i+1} - q_i) \mathcal{D}_{\ell-i-1} \right] \quad (4.7)
 \end{aligned}$$

$$\begin{array}{c}
\text{Diagram: A sequence of nodes connected by dots. Nodes are labeled with 0 or 1 above them. Brackets below group some nodes: } n, p_1, p_{\ell+1}, 2k_{1+1}, p_2, 2k_{\ell'-1+1}, u. \text{ The last node is black.} \\
(4.8)
\end{array}$$

[illegible]

[illegible]

$$\begin{array}{c}
\begin{array}{cc}
1 & 1 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & \frac{1}{2} & 0 & 0 & \frac{1}{2} & \frac{1}{2} & \dots & 0 & 0 & \frac{1}{2} & \frac{1}{2} & 0 & 0 & \frac{1}{2} & \dots & 0 & 0 & \frac{1}{2} & \frac{1}{2} & 0 & 0 & \frac{1}{2} \\
\circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \dots & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \dots & \circ & \circ & \circ & \circ & \circ & \circ & \circ & \bullet \\
\hline
n & & & 2k_1+1 & & & & 2k_2+1 & & & & & & 2k_{\ell-1}+1 & & & & 2k_{\ell}+1 & & & & & & & & & & & &
\end{array} \\
\hline
\underline{2N+1} = \mathcal{D}_{n+\ell} \oplus \mathcal{D}_{\ell-1/2} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1/2}
\end{array} \tag{4.11}$$

4.2 CDD for C_N

We obtain the CDD for C_N by folding A_{2N-1} . A A_N -CDD will give a CDD of C_N iff all the $\mathcal{D}_j$ with $j \in \mathbb{Z}$ have an even multiplicity (in the fundamental of A_{2N-1}).

4.2.1 Integral CDDs

We start again with the CDD (3.4). To get an even multiplicity for $\mathcal{D}_{n+\ell}$ one must impose $n = 0$ and p_1 odd. Then, looking at the other $\mathcal{D}$ -representations leads to the constraints ($n = 0$, $p_i = 2k_i + 1$), and the folded CDD (4.12). The CDD (3.5) can always be folded: we obtain then the CDD (4.13).

$$\begin{array}{c}
\begin{array}{cccccccccccccccccccc}
0 & & 0 & & 1 & & 0 & & 0 & & 1 & & 0 & & 0 & & \cdots & & 1 & & 0 & & 0 & & 1 & & 0 & & 0 & & 0
\end{array} \\
\begin{array}{cccccccccccccccccccc}
\bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \bullet & \cdots & \bullet & \circ
\end{array} \\
\begin{array}{cccccccccccccccccccc}
\underbrace{\hspace{1.5cm}}_{2k_1+1} & & \underbrace{\hspace{1.5cm}}_{2k_2+1} & & \underbrace{\hspace{1.5cm}}_{2k_3+1} & & \cdots & & \underbrace{\hspace{1.5cm}}_{2k_{\ell-1}+1} & & \underbrace{\hspace{1.5cm}}_{k_\ell}
\end{array}
\end{array}
\quad (4.12)$$

[illegible]

4.2.2 Half-integral CDDs

Starting with (3.6), we have to know whether $n + \ell + \ell' - 1$ is even or odd. In the first case, one must impose $n = 0$, so that we have the constraints ($n = 0, \ell' = \ell - 1, p_i = 2k_i + 1$) and obtain (4.15⁻). In the second case, one is left with two possibilities: ($n = 0, \ell' = \ell, q_i = 2k_i + 1$) that leads to (4.14), or ($n = 1, \ell' = \ell - 1, p_i = 2k_i + 1$) that leads to (4.15⁺).

$$\begin{array}{c} \text{Diagram showing a sequence of nodes connected by dots. Nodes are labeled with 0 or } \frac{1}{2}. \text{ Brackets below group nodes into segments labeled } p_1, 2k_1+1, p_2, 2k_2+1, \dots, p_\ell, k_\ell. \\ \underline{2N} = \mathcal{D}_{\ell-1/2} \oplus \mathcal{D}_{\ell-1} \oplus_{i=0}^{\ell-1} [(p_{i+1} - p_i) \mathcal{D}_{\ell-i-1/2} \oplus 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1}] \end{array} \quad (4.14)$$

$$\begin{aligned}
& \left(\overset{\frac{1}{2}}{\bullet} \right) \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{2k_1+1} \overset{\frac{1}{2}}{\bullet} \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{q_1} \overset{\frac{1}{2}}{\bullet} \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{2k_2+1} \overset{\frac{1}{2}}{\bullet} \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{q_2} \cdots \overset{\frac{1}{2}}{\bullet} \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{q_{\ell-1}} \overset{\frac{1}{2}}{\bullet} \underbrace{\overset{0}{\bullet} \cdots \overset{0}{\bullet}}_{k_{\ell}} \bigcirc \\
\underline{2N} = & \begin{cases} \mathcal{D}_{\ell-1/2} \oplus \mathcal{D}_{\ell-1} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1} \oplus_{i=0}^{\ell-2} (q_{i+1} - q_i) \mathcal{D}_{\ell-i-3/2} & \text{with } \bigcirc \\ \mathcal{D}_{\ell-1} \oplus \mathcal{D}_{\ell-3/2} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1} \oplus_{i=0}^{\ell-2} (q_{i+1} - q_i) \mathcal{D}_{\ell-i-3/2} & \text{without } \bigcirc \end{cases}
\end{aligned} \tag{4.15}$$

4.2.3 Mixed CDDs

For the CDD (3.7), and recalling that $n \neq 0$, we have the constraints ($m = 0$, $\ell' = \ell$, $q_i = 2k_i + 1$) to get (4.16), and ($m = 1$, $\ell' = \ell - 1$, $p_i = 2k_i + 1$) to find (4.17).

For the CDD (3.8), $(n = 0, \ell'' = \ell' - 1, p_i = 2k_i + 1)$ leads to (4.18), while $(\ell'' = \ell', q_i = 2k_i + 1)$ produces (4.19).

[illegible]

$$\begin{array}{c}
\text{Diagram: A sequence of nodes connected by dots. Nodes are labeled with weights above them: } 1, \frac{1}{2}, 0, 0, \frac{1}{2}, 0, 0, \frac{1}{2}, 0, 0, \frac{1}{2}, 0, 0, \dots, \frac{1}{2}, 0, 0, \frac{1}{2}, 0, 0, 0. \\
\text{Brackets below group nodes into sets: } n, 2k_1+1, q_1, 2k_2+1, q_2, \dots, q_{\ell-1}, k_\ell. \\
\underline{2N} = \mathcal{D}_{n+\ell-1/2} \oplus \mathcal{D}_{\ell-1} \oplus_{i=0}^{\ell-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1} \oplus_{i=0}^{\ell-2} (q_{i+1} - q_i) \mathcal{D}_{\ell-i-3/2}
\end{array} \quad (4.17)$$

[illegible]

$$\begin{array}{c}
\text{Diagram: A horizontal chain of nodes. Nodes are labeled with values above them: } 1, \dots, 1, 0, 0, 1, \dots, 1, 0, 0, \frac{1}{2}, 0, 0, \frac{1}{2}, \dots, 0, 0, \frac{1}{2}, 0, 0, 0. \\
\text{Braces below the chain group nodes into segments: } n, p_1, \dots, p_{\ell+1}, 2k_1+1, \dots, p_{\ell'+1}, k_{\ell'}. \\
\text{Equation (4.19): } \underline{2N} = \mathcal{D}_{n+\ell'+1/2} \oplus_{i=0}^{\ell+\ell'-1} (p_{i+1} - p_i) \mathcal{D}_{\ell+\ell'-i-1/2} \oplus \mathcal{D}_{\ell'-1} \oplus_{i=0}^{\ell'-1} 2(k_{i+1} - k_i) \mathcal{D}_{\ell-i-1}
\end{array}$$

5 CDDs of D_N

The CDDs of D_N are obtained thanks to the inclusion $D_N \subset B_N$. As the dimension of the fundamental representation of D_N is the one of B_N minus 1, we must select in the CDDs of B_N those which correspond to fundamental representation decomposition with at least one $\mathcal{D}_0$. Then, from the relation between the defining vectors and the CDDs:

$$\left. \begin{array}{l} \text{Diagram 1: } \textcircled{s_1} - \textcircled{s_2} - \dots - \textcircled{s_{n-2}} - \textcircled{s_{n-1}} = \textcircled{s_n} \text{ with } s_i = f_i - f_{i+1}, i = 1, \dots, n-1 \text{ and } s_n = f_n \\ \text{Diagram 2: } \textcircled{s_1} - \textcircled{s_2} - \dots - \textcircled{s_{n-2}} - \textcircled{s_{n-1}} \text{ with } s_i = f_i - f_{i+1}, i = 1, \dots, n-1 \text{ and } s'_n = f_{n-1} + f_n \end{array} \right\} \Rightarrow s'_n = 2s_n + s_{n-1}$$

it is easy to deduce the CDDs of D_N from the CDDs of B_N . We recall that the condition (4.1) is always understood.

As far as the decomposition of the fundamental is concerned, the different cases related to the same CDD of B_N will provide the same sort of decomposition. In fact these decompositions are just the ones of B_N (minus one $\mathcal{D}_0$) so that we do not re-write them in this section.

5.1 Integral CDDs

Let us start with the CDD (4.2). To get a $\mathcal{D}_0$ representation, we must impose $p_\ell = 2u + 2 > p_{\ell-1}$. Then, the computation of s'_n and of the grades entering in the "tail" of the CDD will lead to four different cases: $u \geq 2$, $u = 1$, $u = 0$ and $\ell > 1$, or $u = 0$ and $\ell = 1$. We can suppose in (4.2) that $p_i \neq 0$ since $p_1 = 0$ can be replaced by $n \rightarrow (n+1)$ and $p_i \rightarrow p_{i-1}$ ($i > 1$). If $u \geq 2$, then $s'_n = s_{n-1} = s_{n-2} = 0$ (CDD (5.1)); for $u = 1$, we get $s'_n = s_{n-1} = 0$ but $s_{n-2} = 1$ (CDD (5.2)); finally, $u = 0$ provides $s'_n = s_{n-1} = 1$ and $s_{n-2} = 1$ or 0 , depending on $\ell = 1$ or $\ell > 1$ respectively (CDDs (5.3) and (5.4) respectively). Note that because of the constraints (4.1) and $2u + 2 > p_{\ell-1}$, $u = 0$ leads to $p_i = 1$, while $u = 1$ imposes $p_i \leq 3$.

The CDD (4.3) is of course not allowed, since it has no $\mathcal{D}_0$ representation, but two other integral CDDs (5.5 and 5.6) are provided by mixed CDDs: we write these CDDs here, but explain them in the section dealing with mixed CDDs.

$$\begin{array}{c} \text{Diagram 5.1: } \textcircled{1} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \dots - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{0} - \textcircled{0} \\ \underbrace{\hspace{1.5cm}}_n \quad \underbrace{\hspace{1.5cm}}_{p_1} \quad \underbrace{\hspace{1.5cm}}_{p_2} \quad \dots \quad \underbrace{\hspace{1.5cm}}_{p_{\ell-1}} \quad \underbrace{\hspace{1.5cm}}_{u-2} \quad \textcircled{0} \\ 5 \leq p_{\ell-1} \leq 2u + 1 \end{array} \quad (5.1)$$

$$\begin{array}{c} \text{Diagram 5.2: } \textcircled{1} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{0} - \dots - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} \\ \underbrace{\hspace{1.5cm}}_n \quad \underbrace{\hspace{1.5cm}}_{p_1} \quad \underbrace{\hspace{1.5cm}}_{p_2} \quad \dots \quad \underbrace{\hspace{1.5cm}}_{p_{\ell-2}} \quad \underbrace{\hspace{1.5cm}}_{p_{\ell-1}} \quad \textcircled{1} \\ 0 \leq p_i \leq 3 \end{array} \quad (5.2)$$

$$\begin{array}{c} \text{Diagram 5.3: } \textcircled{1} - \textcircled{1} - \textcircled{1} - \dots - \textcircled{1} - \textcircled{1} - \textcircled{1} \\ \textcircled{1} \end{array} \quad (5.3)$$

$$\begin{array}{c} \text{Diagram 5.4: } \textcircled{1} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \dots - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{1} \\ \underbrace{\hspace{1.5cm}}_n \quad \textcircled{1} \quad (\ell - 1) \text{ nodes } \textcircled{0} \end{array} \quad (5.4)$$

$$\begin{array}{c} \text{Diagram 5.5: } \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \dots - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} - \textcircled{1} - \textcircled{0} - \textcircled{0} \\ \underbrace{\hspace{1.5cm}}_{2k_1+1} \quad \underbrace{\hspace{1.5cm}}_{2k_2+1} \quad \dots \quad \underbrace{\hspace{1.5cm}}_{2k_\ell+1} \quad \underbrace{\hspace{1.5cm}}_{2k_{\ell+1}-1} \quad \textcircled{1} \\ \textcircled{0} \end{array} \quad (5.5)$$

$$\begin{array}{ccccccc}
0 & 1 & 0 & 1 & \dots & 0 & 1 & 0 & \begin{array}{c} \circ 1 \\ | \\ \circ 1 \end{array} & 0
\end{array}
\quad (5.6)$$

5.2 Half-integral CDDs

As for the integral case, one has to distinguish $u = 0$, $u = 1$ and $u \geq 2$. However, as there are two intertwined series (the odd one being never absent), $\ell = 1$ (which should be replaced here by $p_{\ell-1} = 0$) is no more particular with respect to $\ell > 1$ ($p_{\ell-1} > 0$).

Starting from (4.4), one obtains the CDDs (5.7), (5.8), and (5.9) for $u \geq 2$, $u = 1$ and $u = 0$ respectively.

(4.5) can be considered only for the case (4.5⁻) with $\ell = 1$: it produces the integral CDD (5.5) with $\ell = 0$.

The CDD (4.6) produces (5.10), (5.11), and (5.12) in the same way we have expressed for (4.4).

$$\begin{array}{ccccccccccc}
(\frac{1}{2}) & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \dots & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ 0 \\ | \\ \circ 0 \end{array}
\end{array}
\quad (5.7)$$

$\underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{q_1} \quad \underbrace{\quad}_{2k_2+1} \quad \underbrace{\quad}_{q_2} \quad \dots \quad \underbrace{\quad}_{2k_{\ell}+1} \quad \underbrace{\quad}_{u-2}$

$5 \leq q_{\ell-1} \leq 2u + 1$

$$\begin{array}{ccccccccccc}
(\frac{1}{2}) & 0 & 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & \dots & \dots & 0 & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ 0 \\ | \\ \frac{1}{2} \end{array}
\end{array}
\quad (5.8)$$

$\underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{q_1} \quad \underbrace{\quad}_{2k_2+1} \quad \underbrace{\quad}_{q_2} \quad \dots \quad \underbrace{\quad}_{q_{\ell-1}} \quad \underbrace{\quad}_{2k_{\ell}+1}$

$0 \leq q_{\ell-1} \leq 3$

$$\begin{array}{ccccccccccc}
(\frac{1}{2}) & 0 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & \dots & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ \frac{1}{2} \\ | \\ \circ \frac{1}{2} \end{array}
\end{array}
\quad (5.9)$$

$\underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{q_1} \quad \underbrace{\quad}_{2k_2+1} \quad \underbrace{\quad}_{q_2} \quad \dots \quad \underbrace{\quad}_{q_{\ell-1}} \quad \underbrace{\quad}_{2k_{\ell}}$

$0 \leq q_{\ell-1} \leq 1$

$$\begin{array}{ccccccccccc}
0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & \dots & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ 0 \\ | \\ \circ 0 \end{array}
\end{array}
\quad (5.10)$$

$\underbrace{\quad}_{p_1} \quad \underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{p_2} \quad \underbrace{\quad}_{2k_2+1} \quad \dots \quad \underbrace{\quad}_{2k_{\ell-1}+1} \quad \underbrace{\quad}_{u-2}$

$5 \leq p_{\ell-1} \leq 2u + 1$

$$\begin{array}{ccccccccccc}
0 & 0 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 & 0 & \dots & \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ 0 \\ | \\ \frac{1}{2} \end{array}
\end{array}
\quad (5.11)$$

$\underbrace{\quad}_{p_1} \quad \underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{p_2} \quad \underbrace{\quad}_{2k_2+1} \quad \dots \quad \underbrace{\quad}_{p_{\ell-1}} \quad \underbrace{\quad}_{2k_{\ell-1}+1}$

$0 \leq p_{\ell-1} \leq 3$

$$\begin{array}{ccccccccccc}
0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & \dots & \dots & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2} & 0 & 0 & \begin{array}{c} \circ \frac{1}{2} \\ | \\ \circ \frac{1}{2} \end{array}
\end{array}
\quad (5.12)$$

$\underbrace{\quad}_{p_1} \quad \underbrace{\quad}_{2k_1+1} \quad \underbrace{\quad}_{p_2} \quad \underbrace{\quad}_{2k_2+1} \quad \dots \quad \underbrace{\quad}_{p_{\ell-2}} \quad \underbrace{\quad}_{p_{\ell-1}} \quad \underbrace{\quad}_{2k_{\ell-1}}$

$0 \leq p_{\ell-1} \leq 1$

5.3 Mixed CDDs

For the three first mixed CDDs of B_N , the calculation is the same as for the half-integral case. The CDD (4.7) give rise to the CDDs (5.13), (5.14), and (5.15). The CDDs (4.8) and (4.9) produce the CDDs (5.16), (5.17), and (5.18) for the former, and (5.19), (5.20), and (5.21) for the latter.

The CDD (4.10) can contribute only when $\ell' = 1$, which leads to two integral CDDs for D_N . For these CDDs, one has to distinguish if $k_{\ell+1} > 0$ or $k_{\ell+1} = 0$. In the first case, one obtains (5.5), and in the second (5.6).

(4.11) do not contribute at all.

$5 \leq q_{\ell-1} \leq 2u + 1$

$0 \leq q_{\ell-1} \leq 3$

$0 \leq q_{\ell-1} \leq 1$

$5 \leq p_{\ell+\ell'-1} \leq 2u + 1$

$0 \leq p_{\ell+\ell'-1} \leq 3$

$0 \leq p_{\ell+\ell'-1} \leq 1$

[illegible]

$$\begin{array}{c}
\textcircled{0} \cdots \textcircled{0} \textcircled{1} \textcircled{0} \cdots \textcircled{0} \cdots \textcircled{1} \textcircled{0} \cdots \textcircled{0} \overset{\frac{1}{2}}{\textcircled{}} \textcircled{0} \textcircled{0} \textcircled{0} \overset{\frac{1}{2}}{\textcircled{}} \textcircled{0} \cdots \textcircled{0} \overset{\frac{1}{2}}{\textcircled{}} \textcircled{0} \textcircled{0} \textcircled{0} \cdots \overset{\frac{1}{2}}{\textcircled{}} \textcircled{0} \textcircled{0} \textcircled{0} \overset{\frac{1}{2}}{\textcircled{}} \textcircled{0} \cdots \textcircled{0} \textcircled{\scriptsize 0} \\
\underbrace{\hspace{1cm}}_{2k_1+1} \quad \underbrace{\hspace{1cm}}_{2k_2+1} \quad \cdots \quad \underbrace{\hspace{1cm}}_{2k_{\ell'+1}+1} \quad \underbrace{\hspace{1cm}}_{q_1} \quad \underbrace{\hspace{1cm}}_{2k_{\ell'+2}+1} \quad \underbrace{\hspace{1cm}}_{q_2} \quad \cdots \quad \underbrace{\hspace{1cm}}_{q_{\ell'-1}} \quad \underbrace{\hspace{1cm}}_{2k_{\ell'+\ell'}+1} \quad \textcircled{\scriptsize 0} \\
0 \leq q_{\ell'-1} \leq 3
\end{array}
\tag{5.20}$$

[illegible]

5.4 Folding of the CDDs of D_N

In the same way that we have folded A_M to get B_N and C_N , it is possible, using the $\mathbb{Z}_2$ symmetry of the "tail" of D_N , to fold it and get B_{N-1} . Note however that the CDDs of D_N do not possess *a priori* the symmetry that allows the folding, so that we have to select the subset of symmetric CDDs, that is to discard the CDDs (5.5) and (5.6). Moreover, the fundamental representation $\underline{2N-1}$ of B_{N-1} is obtained from the $\underline{2N}$ representation of D_N by $\underline{2N-1} = \underline{2N} - \mathcal{D}_0$: we must also select the CDDs that correspond to a decomposition with at least one representation $\mathcal{D}_0$. This prescription will suppress all the CDDs associated with $u = 0$. Then, one can fold the remaining CDDs of D_N , and re-find the results given in section 4.1, the cases $u = 1$ becoming a "real" case of $u \geq 2$.

As an exercise, let us also compute the CDDs of G_2 thanks to the $\mathbb{Z}_3$ symmetry of D_4 . G_2 -CDDs are obtained from D_4 -CDDs by identifying its three external nodes. We have also to impose that the decomposition of the $\mathfrak{g}$ of D_4 possesses a $\mathcal{D}_0$ representation, since the fundamental of G_2 has dimension 7. The CDDs of D_4 are:

(5.22)

The first line corresponds to non $\mathbb{Z}_3$ -symmetric CDDs. Note however that this line shows how the "new" outer isomorphism (induced by the enlarged symmetry of D_4 w.r.t. D_N , $N \neq 4$) relates

some of the $Sl(2)$ of D_4 . On the second line lies the only CDD of D_4 which is $\mathbb{Z}_3$ symmetric but which does not possess a $\mathcal{D}_0$ representation in its fundamental representation. Thus, we fold the CDDs of the last line and get the CDDs of G_2 :

$$\begin{array}{cccc} \overset{1}{\bullet} \equiv \overset{1}{\circ} & \overset{0}{\bullet} \equiv \overset{1}{\circ} & \overset{0}{\bullet} \equiv \overset{\frac{1}{2}}{\circ} & \overset{\frac{1}{2}}{\bullet} \equiv \overset{0}{\circ} \end{array} \quad (5.23)$$

6 Application to W algebras

Most (and perhaps all) the W algebras can be realized in the context of non Abelian Toda theories. Such models are deduced from a WZW action $S_0(g)$, based on a group G of Lie algebra $\mathcal{G}$, by imposing some constraints on the (chiral) Kac-Moody currents of $S_0(g)$. These constraints are themselves induced by a gradation H of $\mathcal{G}$. Then a Lagrange multiplier treatment and a Gauss decomposition $g = g_+ g_0 g_-$ lead to a new action $S(g_0)$ that depends only on the 0-grade (w.r.t. H) component g_0 . This provides an easy-to-handle framework for the realization of W algebras (which are symmetry of non Abelian Toda theories) in term of the Kac-Moody generators on the one hand, and in terms of free fields on the other hand. The W algebra obtained is naturally associated to an $Sl(2)$ $\mathcal{G}$ -subalgebra, H being the Cartan generator of this $Sl(2)$ subalgebra. In fact, the $Sl(2)$ embeddings in $\mathcal{G}$ classify the different W algebras which can be obtained by such a procedure. These algebras are denoted $\mathcal{W}(\mathcal{G}, \mathcal{K})$, where $\mathcal{K}$ is the $\mathcal{G}$ -subalgebra whose principal $Sl(2)$ is the $Sl(2)$ algebra under consideration. Since these W algebras are classified by the different $Sl(2)$ in $\mathcal{G}$, it is natural to relate also these algebras to CDDs. Let us immediately remark that such an approach will be particularly useful for the folding of W algebras [5], since the CDDs of B_N and C_N already have been construct using this method. We give below other examples of the use of CDDs in the study of $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras.

6.1 Characterization of $\mathcal{W}(\mathcal{G}, \mathcal{K})$

As explain above, since the $\mathcal{W}(\mathcal{G}, \mathcal{K})$ are classified by $Sl(2)$ subalgebras in $\mathcal{G}$, we can think of the $\mathcal{G}$ -CDDs as the Dynkin Diagrams for the $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras. We want here to show how the use of the CDDs, combined with the results obtained in [2] give a lot of informations on the structure of $\mathcal{W}(\mathcal{G}, \mathcal{K})$ with a minimal of calculation. Using only the decomposition of the fundamental $\mathcal{G}$ -representation with respect to this $Sl(2)$, [2] were able to compute the spin contents of all the $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras. Moreover, extending the $Sl(2)$ algebra to an $Sl(2) \oplus U(1)$ algebra, they also give the eigenvalues of all the W generators with respect to some special $U(1)$. From this work, and the knowledge of the fundamental representation decomposition, it is possible to make contact between CDDs and $\mathcal{W}(\mathcal{G}, \mathcal{K})$ algebras. Moreover, the knowledge of the grades of the simple roots allows us to compute the Kac-Moody subalgebra which is in $\mathcal{W}(\mathcal{G}, \mathcal{K})$, up to $U(1)$ factors. In fact, this Kac-Moody algebra is just the "affinization" of the algebra one obtains by removing in the CDD all the nodes that have not a 0-grade. Note that the $U(1)$ factors that the CDD cannot determine are just the ones whose eigenvalues are computed in [2], so that combining the two approaches gives almost all the informations on $\mathcal{W}(\mathcal{G}, \mathcal{K})$. The results are given in the tables below. We will use the notations

$$\begin{aligned} r_i &= p_{i+1} - p_i \quad i \geq 1 & \text{and} \quad r_0 &= p_1 \\ s_i &= q_{i+1} - q_i \quad i \geq 1 & \text{and} \quad s_0 &= q_1 \\ l_i &= k_{i+1} - k_i \quad i \geq 1 \end{aligned} \quad (6.1)$$

6.1.1 $\mathcal{W}(A_N, \mathcal{K})$ algebras

The only regular subalgebras of A_N are (sum of) A_M subalgebras. When reducing the fundamental representation with respect to the principal $Sl(2)$ of A_M , a $\mathcal{D}_{M/2}$ representation will appear. Thus, we identify the form of $\mathcal{K}$ thanks to the rule

$$\mathcal{D}_{p/2} \text{ in fundamental} \longrightarrow A_p \text{ in } \mathcal{K} \quad (6.2)$$

Looking at all the CDDs of A_N we obtain the following table

CDD	$\mathcal{K}$ in $\mathcal{W}(A_N, \mathcal{K})$	KM subalg. in $\mathcal{W}$ (up to $U(1)$ factors)
(3.4)	$A_{2n+2\ell-2} \oplus_{i=0}^{\ell-2} r_i A_{2\ell-2i-2}$	$A_{p_\ell} \oplus_{i=1}^{\ell-1} 2A_{p_i}$
(3.5)	$A_{2n+2\ell-1} \oplus_{i=0}^{\ell-2} r_i A_{2\ell-2i-1}$	$2 \oplus_{i=1}^{\ell} A_{p_i}$
(3.6)	$A_{n+\ell+\ell'-1} \oplus A_{n+\ell+\ell'-2} \oplus$ $\oplus_{i=0}^{\ell-1} r_i A_{\ell+\ell'-2i-1} \oplus_{i=0}^{\ell'-1} s_i A_{\ell+\ell'-2i-2}$	$2 \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{q_i}] \oplus A_{p_\ell}$ (if $\ell' = \ell - 1$) $2 \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{q_i}] \oplus 2A_{p_\ell} \oplus A_{q_\ell}$ (if $\ell' = \ell$)
(3.7) ($n \neq 0$)	$A_{2n+m+\ell+\ell'-1} \oplus A_{m+\ell+\ell'-2} \oplus$ $\oplus_{i=0}^{\ell-1} r_i A_{\ell+\ell'-2i-1} \oplus_{i=0}^{\ell'-1} s_i A_{\ell+\ell'-2i-2}$	$2 \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{q_i}] \oplus A_{p_\ell}$ (if $\ell' = \ell - 1$) $2 \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{q_i}] \oplus 2A_{p_\ell} \oplus A_{q_\ell}$ (if $\ell' = \ell$)
(3.8) ($n + \ell \neq 0$)	$A_{2n+2\ell+\ell'+\ell''-1} \oplus_{i=0}^{\ell+\ell'-1} r_i A_{2\ell+\ell'+\ell''-2i-1}$ $\oplus A_{\ell+\ell'-2} \oplus_{i=0}^{\ell''-1} s_i A_{\ell+\ell'-2i-2}$	$2 [\oplus_{i=1}^{\ell+\ell'-1} A_{p_i} \oplus_{i=1}^{\ell''-1} A_{q_i}] \oplus A_{p_{\ell+\ell'}}$ (if $\ell'' = \ell' - 1$) $2 \oplus_{i=1}^{\ell+\ell'-1} [A_{p_i} \oplus A_{q_i}] \oplus 2A_{p_{\ell+\ell'}} \oplus A_{q_{\ell''}}$ (if $\ell'' = \ell'$)

(6.3)

The spin contents and the hypercharges are then easily computed from the decompositions given in table 3.9, following the scheme developed in [2].

6.1.2 $\mathcal{W}(B_N, \mathcal{K})$ algebras

B_N contains as regular subalgebras A_M , B_M and D_M . The relation between these subalgebras and the decomposition of the fundamental with respect to their principal $Sl(2)$ have been given in [2]: each A_M contributes to the fundamental decomposition by a term $2\mathcal{D}_{M/2}$, and B_M or D_M provide a $\mathcal{D}_M$ representation (we use the notation B_1 for the A_1 algebra of index 2, constructed on a short root). Moreover, in B_N , the principal $Sl(2)$ constructed on a D_M algebra is the same as the one constructed on a B_M algebra, so that we can neglect these inclusions. Thus, the rules are in this case

$$\begin{cases} \mathcal{D}_p \longrightarrow B_p \\ 2\mathcal{D}_{p/2} \longrightarrow A_{p-1} \end{cases} \quad (6.4)$$

CDD	$\mathcal{K}$ in $\mathcal{W}(B_N, \mathcal{K})$	KM subalg. in $\mathcal{W}$ (up to $U(1)$ factors)
(4.2)	$B_{n+\ell-1} \oplus p_1 B_{\ell-1} \oplus_{i=1}^{\ell-2} r_i B_{\ell-i-1}$	$B_{u+1} \oplus_{i=1}^{\ell-1} A_{p_i}$
(4.3)	B_N	$\emptyset$
(4.4 ⁻)	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)B_{\ell-1} \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i B_{\ell-i-1}$	$B_{u+1} \oplus A_{2k_\ell+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(4.4 ⁺)	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)B_\ell \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i B_{\ell-i-1}$	$B_{u+1} \oplus A_{2k_\ell+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(4.5 ⁻)	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)B_{\ell-1} \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1}$	$\oplus_{i=1}^{\ell} A_{2k_i+1}$
(4.5 ⁺)	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)B_\ell \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1}$	$\oplus_{i=1}^{\ell} A_{2k_i+1}$
(4.6)	$(p_1 + 1)B_{\ell-1} \oplus (k_1 + 1)A_{2\ell-3} \oplus$ $\oplus_{i=1}^{\ell-2} [r_i B_{\ell-i-1} \oplus l_i A_{2\ell-2i-3}]$	$B_{u+1} \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{2k_i+1}]$
(4.7)	$B_{n+\ell} \oplus (k_1 + 1)A_{2\ell-1} \oplus q_1 B_\ell \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i B_{\ell-i-1}$	$A_{2k_\ell+1} \oplus B_{u+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(4.8)	$B_{n+\ell+\ell'-1} \oplus q_1 B_{\ell+\ell'-1} \oplus_{i=1}^{\ell+\ell'-2} r_i B_{\ell+\ell'-i-1}$ $\oplus (k_1 + 1)A_{2\ell'-3} \oplus_{i=0}^{\ell'-2} l_i A_{2\ell'-2i-3}$	$B_{u+1} \oplus_{i=1}^{\ell+\ell'-1} A_{p_i} \oplus_{i=1}^{\ell'-1} A_{2k_i+1}$
(4.9)	$(k_1 + 1)A_{2\ell+2\ell'-1} \oplus_{i=1}^{\ell+\ell'-1} l_i A_{2\ell+2\ell'-2i-1} \oplus$ $\oplus (q_1 + 1)B_{\ell'-1} \oplus_{i=1}^{\ell'-2} s_i B_{\ell'-i-1}$	$B_{u+1} \oplus_{i=1}^{\ell+\ell'} A_{2k_i+1} \oplus_{i=1}^{\ell'-1} A_{q_i}$
(4.10)	$(k_1 + 1)A_{2\ell+2\ell'-1} \oplus B_{\ell'-1} \oplus$ $\oplus_{i=1}^{\ell+\ell'-1} l_i A_{2\ell+2\ell'-2i-1}$	$\oplus_{i=1}^{\ell+\ell'} A_{2k_i+1}$
(4.11)	$B_{n+\ell} \oplus (k_1 + 1)A_{2\ell-1} \oplus_{i=1}^{\ell+\ell'-1} l_i A_{2\ell-2i-1}$	$\oplus_{i=1}^{\ell} A_{2k_i+1}$

(6.5)

6.1.3 $\mathcal{W}(C_N, \mathcal{K})$ algebras

As in the previous case, we can write C_1 to distinguish the index 1 A_1 algebra constructed on a long root, from the index 2 A_M subalgebras. Then, the correspondence between $\mathcal{D}_M$ representations and (regular) subalgebras will be:

$$\begin{cases} \mathcal{D}_{p-1/2} \longrightarrow C_p \\ 2\mathcal{D}_{p/2} \longrightarrow A_p \end{cases} \quad (6.6)$$

CDD	$\mathcal{K}$ in $\mathcal{W}(C_N, \mathcal{K})$	KM subalg. in $\mathcal{W}$ (up to $U(1)$ factors)
(4.12)	$(k_1 + 1)A_{2\ell-2} \oplus_{i=1}^{\ell-2} l_i A_{2\ell-2i-2}$	$C_{k_\ell+1} \oplus_{i=1}^{\ell-1} A_{2k_i+1}$
(4.13)	$(p_1 + 1)C_{n+\ell} \oplus_{i=1}^{\ell-2} r_i C_{\ell-i}$	$\oplus_{i=1}^{\ell} A_{p_i}$
(4.14)	$(p_1 + 1)C_\ell \oplus (k_1 + 1)A_{2\ell-2} \oplus$ $\oplus_{i=1}^{\ell-2} [r_i C_{\ell-i} \oplus l_i A_{2\ell-2i-2}]$	$\oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{2k_i+1}]$ $\oplus A_{p_\ell} \oplus C_{k_\ell+1}$
(4.15 ⁻)	$(k_1 + 1)A_{2\ell-2} \oplus (q_1 + 1)C_{\ell-1} \oplus$ $\oplus_{i=1}^{\ell-2} [l_i A_{2\ell-2i-2} \oplus s_i C_{\ell-i-1}]$	$\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}] \oplus C_{k_\ell+1}$
(4.15 ⁺)	$(k_1 + 1)A_{2\ell-2} \oplus (q_1 + 1)C_\ell \oplus$ $\oplus_{i=1}^{\ell-2} [l_i A_{2\ell-2i-2} \oplus s_i C_{\ell-i-1}]$	$\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}] \oplus C_{k_\ell+1}$
(4.16)	$C_{n+\ell} \oplus (k_1 + 1)A_{2\ell-2} \oplus q_1 C_\ell \oplus$ $\oplus_{i=1}^{\ell-2} [r_i C_{\ell-i-1} \oplus l_i A_{2\ell-2i-2}]$	$\oplus_{i=1}^{\ell} A_{p_i} \oplus_{i=1}^{\ell-1} A_{2k_i+1} \oplus C_{k_\ell+1}$
(4.17)	$C_{n+\ell} \oplus (k_1 + 1)A_{2\ell-2} \oplus$ $\oplus_{i=1}^{\ell-2} [l_i A_{2\ell-2i-2} \oplus s_i C_{\ell-i-1}]$	$\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}] \oplus C_{k_\ell+1}$
(4.18)	$(k_1 + 1)A_{2\ell+2\ell'-2} \oplus_{i=1}^{\ell+\ell'-2} l_i A_{2\ell+2\ell'-2i-2} \oplus$ $\oplus (q_1 + 1)C_{\ell'-1} \oplus_{i=1}^{\ell'-2} s_i C_{\ell'-i-1}$	$\oplus_{i=1}^{\ell+\ell'-1} A_{2k_i+1} \oplus_{i=1}^{\ell'-1} A_{q_i} \oplus C_{k_{\ell+\ell'}+1}$
(4.19)	$C_{n+\ell+\ell'} \oplus p_1 C_{\ell+\ell'} \oplus_{i=1}^{\ell+\ell'-2} r_i C_{\ell+\ell'-i} \oplus$ $\oplus (k_1 + 1)A_{2\ell'-2} \oplus_{i=1}^{\ell'-2} l_i A_{2\ell'-2i-2}$	$\oplus_{i=1}^{\ell+\ell'} A_{p_i} \oplus_{i=1}^{\ell'-1} A_{2k_i+1} \oplus C_{k_{\ell'}+1}$

(6.7)

6.1.4 $\mathcal{W}(D_N, \mathcal{K})$ algebras

As far as D_N algebra are concerned, one can proceed in the same way as for B_N (replacing B_M by D_M) except for the irregular embeddings $B_k \oplus B_{N-k} \subset D_N$, that cannot be replaced by D_M embeddings. It is natural to think of them as related to the particular cases $u = 0$ or $u = 1$ of the integral CDDs. As these irregular embeddings should not survive to the folding $D_N \rightarrow B_{N-1}$, we focus on $u = 0$ and $\ell > 1$ ($\ell = 1$ is the principal CDD). Then, looking at the $Sl(2)$ decomposition of the CDD (5.4), we obtain $\underline{2N} = \mathcal{D}_{n+\ell-1} \oplus \mathcal{D}_{\ell-1}$ which indeed corresponds to the reduction with respect to the principal $Sl(2)$ of $^2 B_{\ell-1} \oplus B_{N-\ell+1}$. Note that the case $\ell = 1$ could also be included in the exceptions (with the convention $B_0 = \emptyset$), since the $Sl(2)$ -principal embedding of B_{N-1} in D_N is also the principal $Sl(2)$ of D_N . Thus the rules are

$$\text{CDDs (5.3) and (5.4)} \longrightarrow B_{n+\ell-1} \oplus B_{\ell-1} \quad (6.8)$$

²We have used the equality $2N = 2n + 2\ell - 1 + 2\ell - 1$, resulting from the decomposition of the fundamental.

$$\text{other CDDs } \left\{ \begin{array}{l} \mathcal{D}_p \longrightarrow D_p \\ \mathcal{D}_{p/2} \longrightarrow A_p \end{array} \right. \quad (6.9)$$

For the three D_N -CDDs ($u = 0$, $u = 1$, $u \geq 2$) associated to one B_N -CDD, the decomposition of the fundamental representation has the same form, so that we will always (up to the above exceptions) get the same type of subalgebra $\mathcal{K}$. However, because the tails have different grades the $\mathcal{G}_0$ subalgebras are different. In the case where $u = 1$, we get two A_1 subalgebras from the tail, instead of the D_{u+1} subalgebra arising when $u \geq 2$. But for $u = 1$, the algebra $D_{u+1} = D_2$ is isomorphic to $2A_1$, so that we include $u = 1$ in the general treatment $u \geq 2$. Note that once more, it is the case $u = 0$ that appears much different from the other ones.

When $u = 0$, we call i_0 the only index such that $p_{i_0} = 0$ and $p_{i_0+1} = 1$ (or $q_{i_0} = 0$ and $q_{i_0+1} = 1$): it is the only index such that $r_{i_0} = 1 \neq 0$ (or $s_{i_0} = 1 \neq 0$). In the case of integral grading we have $i_0 = \ell - 1$.

CDD	$\mathcal{K}$ in $\mathcal{W}(D_N, \mathcal{K})$	KM subalg. in $\mathcal{W}$ (up to $U(1)$ factors)
$\left. \begin{array}{l} (5.1) \\ (5.2) \end{array} \right\}$	$D_{n+\ell-1} \oplus p_1 D_{\ell-1} \oplus \bigoplus_{i=1}^{\ell-2} r_i D_{\ell-i-1}$	$D_{u+1} \oplus \bigoplus_{i=1}^{\ell-1} A_{p_i}$
$\left. \begin{array}{l} (5.3) \\ (5.4) \end{array} \right\}$	$B_{N-\ell+1} \oplus B_{\ell-1}$	$(\ell-1)A_{p_{\ell-1}}$ with $p_{\ell-1} = \begin{cases} 1 \\ 0 \end{cases}$
(5.5)	$(k_1+1)A_{2\ell+1} \oplus \bigoplus_{i=1}^{\ell} l_i A_{2\ell-2i+1}$	$\bigoplus_{i=1}^{\ell+1} A_{2k_\ell+1}$
(5.6)	$(k_1+1)A_1$	ℓA_1

(6.10)

CDD	$\mathcal{K}$ in $\mathcal{W}(D_N, \mathcal{K})$	KM subalg. in $\mathcal{W}$ (up to $U(1)$ factors)
$\left. \begin{array}{l} (5.7^+) \\ (5.8^+) \end{array} \right\}$	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)D_\ell \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i D_{\ell-i-1}$	$D_{u+1} \oplus A_{2k_\ell+1} \oplus$ $\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
$\left. \begin{array}{l} (5.7^-) \\ (5.8^-) \end{array} \right\}$	$(k_1 + 1)A_{2\ell-1} \oplus (q_1 + 1)D_{\ell-1} \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i D_{\ell-i-1}$	$D_{u+1} \oplus A_{2k_\ell+1} \oplus$ $\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(5.9 ⁻)	$(k_1 + 1)A_{2\ell-1} \oplus D_{\ell-1} \oplus$ $\oplus D_{\ell-i_0-1} \oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1}$	$A_{2k_\ell+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(5.9 ⁺)	$(k_1 + 1)A_{2\ell-1} \oplus D_\ell \oplus$ $\oplus D_{\ell-i_0-1} \oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1}$	$A_{2k_\ell+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
$\left. \begin{array}{l} (5.10) \\ (5.11) \end{array} \right\}$	$(p_1 + 1)D_\ell \oplus (k_1 + 1)A_{2\ell-3} \oplus$ $\oplus_{i=1}^{\ell-2} [r_i D_{\ell-i-1} \oplus l_i A_{2\ell-2i-3}]$	$D_{u+1} \oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{2k_i+1}]$
(5.12)	$D_\ell \oplus (k_1 + 1)A_{2\ell-3} \oplus$ $\oplus D_{\ell-i_0-1} \oplus_{i=1}^{\ell-2} l_i A_{2\ell-2i-3}$	$\oplus_{i=1}^{\ell-1} [A_{p_i} \oplus A_{2k_i+1}]$
$\left. \begin{array}{l} (5.13) \\ (5.14) \end{array} \right\}$	$(k_1 + 1)A_{2\ell-1} \oplus D_{n+\ell} \oplus q_1 D_\ell \oplus$ $\oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1} \oplus_{i=1}^{\ell-2} s_i D_{\ell-i-1}$	$D_{u+1} \oplus A_{2k_\ell+1} \oplus$ $\oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
(5.15)	$(k_1 + 1)A_{2\ell-1} \oplus D_{n+\ell} \oplus$ $\oplus D_{\ell-i_0-1} \oplus_{i=1}^{\ell-1} l_i A_{2\ell-2i-1}$	$A_{2k_\ell+1} \oplus_{i=1}^{\ell-1} [A_{2k_i+1} \oplus A_{q_i}]$
$\left. \begin{array}{l} (5.16) \\ (5.17) \end{array} \right\}$	$D_{n+\ell+\ell'-1} \oplus_{i=1}^{\ell+\ell'-2} r_i D_{\ell+\ell'-i-1} \oplus p_1 D_{\ell+\ell'-1}$ $\oplus (k_1 + 1)A_{2\ell'-3} \oplus_{i=1}^{\ell'-2} l_i A_{2\ell'-2i-3}$	$D_{u+1} \oplus_{i=1}^{\ell+\ell'-1} A_{p_i} \oplus_{i=1}^{\ell'} A_{2k_i+1}$
(5.18)	$D_{n+\ell+\ell'-1} \oplus D_{\ell+\ell'-i_0-1} \oplus$ $\oplus (k_1 + 1)A_{2\ell'-3} \oplus_{i=1}^{\ell'-2} l_i A_{2\ell'-2i-3}$	$\oplus_{i=1}^{\ell+\ell'-1} A_{p_i} \oplus_{i=1}^{\ell'} A_{2k_i+1}$
$\left. \begin{array}{l} (5.19) \\ (5.20) \end{array} \right\}$	$(k_1 + 1)A_{2\ell+2\ell'-1} \oplus (q_1 + 1)D_{\ell'-1} \oplus$ $\oplus_{i=1}^{\ell'-2} D_{\ell'-i-1} \oplus_{i=1}^{\ell+\ell'-1} l_i A_{2\ell+2\ell'-2i-1}$	$D_{u+1} \oplus_{i=1}^{\ell'-1} A_{q_i} \oplus_{i=1}^{\ell+\ell'} A_{2k_i+1}$
(5.21)	$(k_1 + 1)A_{2\ell+2\ell'-1} \oplus D_{\ell'-1} \oplus D_{\ell'-i_0-1}$ $\oplus_{i=1}^{\ell+\ell'-1} l_i A_{2\ell+2\ell'-2i-1}$	$\oplus_{i=1}^{\ell'-1} A_{q_i} \oplus_{i=1}^{\ell+\ell'} A_{2k_i+1}$

(6.11)

6.2 Duality between B_N and C_N

It has been proven in [6] that the $\mathcal{W}(B_N, B_N)$ and $\mathcal{W}(C_N, C_N)$ exhibit a duality, in the sense that the Abelian Toda model constructed on C_N is obtained from the Abelian Toda model built on B_N by replacing the roots system by its dual, and the coupling constant β by $\frac{-1}{\beta}$.

Using the CDDs, it is possible to see which of the (other) $\mathcal{W}(B_N, \mathcal{K})$ and $\mathcal{W}(C_N, \mathcal{K})$ algebras can be dual. Indeed, using the duality between B_N and C_N DDs that consists in exchanging the colors black and white in the nodes, one obtains the following candidates for the "B-C duality"

$$\begin{array}{c}
 \textcircled{0} - \underbrace{\textcircled{0}}_{2k_1+1} - \textcircled{1} - \underbrace{\textcircled{0}}_{2k_2+1} - \textcircled{1} - \underbrace{\textcircled{0}}_{2k_3+1} - \dots - \underbrace{\textcircled{1}}_{2k_{\ell-1}+1} - \underbrace{\textcircled{0}}_{k_\ell} \\
 \end{array} \quad (6.12)$$

$$\begin{array}{c} 1 \quad 1 \quad \quad 1 \quad 1 \quad 1 \\ \circ - \circ - \cdots - \circ - \circ = \circ \end{array} \quad (6.13)$$

$$\begin{array}{ccccccc}
0 & \dots & 0 & \frac{1}{2} & 0 & \dots & 0 \\
\circ & \text{---} & \circ & \circ & \circ & \text{---} & \circ \\
& \underbrace{\hspace{1cm}}_{2k_1+1} & & \underbrace{\hspace{1cm}}_{2j_1+1} & & \underbrace{\hspace{1cm}}_{2k_2+1} & \underbrace{\hspace{1cm}}_{2j_2+1} \dots \underbrace{\hspace{1cm}}_{2k_\ell+1} \underbrace{\hspace{1cm}}_{j_\ell}
\end{array}
\quad (6.14)$$

$$\begin{array}{c}
0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad 0 \\
\circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \quad \circ \\
\hline
2k_1+1 \quad 2j_1+1 \quad 2k_2+1 \quad 2j_2+1 \quad \cdots \quad 2j_{\ell-1}+1 \quad k_\ell
\end{array} \quad (6.15)$$

[illegible]

$$\begin{array}{c}
1 \quad 1 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad \frac{1}{2} \quad 0 \quad 0 \quad 0 \quad 0 \\
\circ \quad \circ \\
\vdots \quad \vdots \\
n \quad 2k_1+1 \quad 2j_1+1 \quad 2k_2+1 \quad 2j_2+1 \quad \dots \quad 2k_\ell+1 \quad j_\ell
\end{array} \quad (6.17)$$

[illegible]

Notice that one has to pay attention, when comparing B_N and C_N CDDs, to the difference between $\ell' = \ell$ and $\ell' = \ell - 1$. For instance, (4.7) seems very similar to (4.17), but has one piece of length q more than the other: to get (6.17), one has to take (4.7) with q_i odd ($i \neq \ell$) on the one hand, but (4.19) with $\ell = 0$, p_i odd ($i \geq 2$) and $p_1 = 0$ on the other hand.

Looking at the fundamental representation, it is easy to see that in most of the cases we have the same decomposition for the B_N and C_N algebras (except one $\mathcal{D}_0$ that allows to go from $\underline{2N}$ to $\underline{2N+1}$). This means that the spin contents of the corresponding W algebras will be different, since it is obtained by a symmetric (antisymmetric) product of the fundamental by its contragredient for C_N (B_N). The exceptions that have different fundamental decomposition for B_N and C_N are (6.13) and (6.17). The principal CDD (6.13) lead to the well-known algebras $\mathcal{W}(B_N, B_N)$ and $\mathcal{W}(C_N, C_N)$, and a direct calculation of the adjoint decomposition for (6.17) shows that the spin content is still different.

Thus, the only W algebras related by a "B-C duality" are $\mathcal{W}(B_N, B_N)$ and $\mathcal{W}(C_N, C_N)$. It is nevertheless possible that the W algebras pointed out in this paragraph are related through another relation weaker than this duality.

7 Conclusion

In this paper, we have presented a classification of CDDs for A,B,C,D algebras, and connected them to the classification of $\mathcal{W}(\mathcal{G},\mathcal{K})$ algebras. The CDD may be viewed as a kind of Dynkin diagram for these algebras, so that one can wonder whether it is still true for a more general class of W algebras. For such a purpose, one can think of enlarging the definition of CDDs by allowing the grades to take any values and try to connect W algebras with "generalized" CDDs in a different way than the $Sl(2)$ classification used here. If it is possible, this will imply that a "generalised" CDD encode all the relevant informations concerning the W algebras (as it is the case for CDDs with $\mathcal{W}(\mathcal{G},\mathcal{K})$ algebras). Note however that a generalised CDD will always be related to a Cartan generator of $\mathcal{G}$ (but not of a $Sl(2)$ $\mathcal{G}$ -subalgebra), through the grades of the simple roots.

Finally, let us remark that a similar treatment for super- W algebras, with some "super-CDDs" classifying $OSp(1|2)$ sub-superalgebras, may perhaps be done. One has first to know if the $OSp(1|2)$ embeddings can be classify by such diagrams: to our knowledge, this has not yet been studied, but the notion of super-defining vectors has already being introduced in [2].

Acknowledgements

I would like to thank Jens Ole Madsen, Jens Lyng Petersen and Paul Sorba for reading this manuscript.

References

- [1] For a general review on W algebras, see P. Bouwknegt and K. Schoutens, preprint CERN-TH-6583 (july 92, to be published in Phys. Reports) and references therein.
- [2] L. Frappat, E. Ragoucy and P. Sorba, "*W-algebras and superalgebras from constrained WZW models: a group theoretical classification.*", preprint ENSLAPP-A-391/92.
- [3] For a general review on non Abelian Toda theories, see L. Fehér, L. O'Raifeartaigh, P. Ruelle, I. Tsutsui and A. Wipf, Phys. Report **222** (dec. 92) *n*^o 1, and references therein.
- [4] E. Dynkin, Amer. Math. Soc. Transl. Ser. 2, **6** (1957) 111.
- [5] L. Frappat, E. Ragoucy and P. Sorba, "*Folding the W-algebras.*", preprint ENSLAPP-A-408/92.
- [6] H.G. Kausch and G.M.T. Watts, preprint Imperial/TH/91-92/17.